



Cerca nel sito Cerca

HOME > GRUPPI DI STUDIO > Testa collo

Prostata

Iort

Gastrointestinale

Mammella

Brachiterapia

Rt metabolica

Cure palliative

Neoplasie cerebrali

Testa collo

Rt stereotassica

Polmone



Associazione Italiana Radioterapia Oncologica



Linee guida multidisciplinari

METASTASI LINFONODALI DEL COLLO A SEDE PRIMITIVA SCOSCIUTA


TUMORI Algoritmi ver

• in nero: gli algoritmi di
Guida NCCN

• In rosso:

Anamnesi
Ispezione
testa e collo e cute
Visita clinica
con fibroscopia

METASTASI LINFONODALI DEL COLLO A SEDE PRIMITIVA SCOSCIUTA



Coordinatore Prof. Corvò

Technical Guidelines of IMRT for Head and Neck Cancer

on behalf of the task group of the Italian Association of Radiation Oncology

Anna Merlotti¹, Daniela Alterio², Riccardo Vigna Taglianti³, Alessandro Muraglia⁴, Luciana Lastrucci⁵, Francesco Deodato, Roberto Manzo⁶, Giuseppina Gambaro⁷, Orietta Caspiani⁸, Francesco Miccicchè⁹, Renzo Corvò¹⁰, Elvio Russi³, Giuseppe Sanguineti¹¹

Neck metastases from unknown primary (cT0)

General remarks

1. Metastatic cervical adenopathy of unknown primary origin represents a very heterogeneous entity. Squamous cell carcinoma is the more frequent histological type, followed by adenocarcinoma and undifferentiated carcinoma. If melanoma, lymphoma, adenocarcinoma (usually chest but also thyroid and salivary) treat accordingly; here we consider only SCC or undifferentiated carcinoma;
2. Some associations between the location of the node and the primary site: superficial parotid: skin; level Ib: oral cavity; level II-Va: NPC; level IV: chest (or abdomen)
3. Work up: